



Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023

Acute Respiratory Infection (ARI)

Covid-19, Influenza and other respiratory viruses

High activity in general practice





In mainland France, last week (2023w06), the incidence rate of ARI cases consulting in general practice was estimated at **284 cases per 100,000 inhabitants (95% CI [260 ; 308])**. This rate is **increasing** compared to week 2023w05 (consolidated data: 231 [213 ; 249]).

- Additional IRA data: page 2
- Influenza data: page 3
- Covid-19 data: page 4
- RSV data: page 5
- Complete national and regional data: page 6

ARI are caused by a variety of respiratory viruses including SARS-CoV-2 (Covid-19), influenza viruses, and other respiratory viruses such as RSV, rhinovirus and metapneumovirus. The purpose of ARI surveillance is to monitor outbreaks of these virus.

Acute diarrhea

Moderate to high activity in general practice



In mainland France, last week (2023w06), the incidence rate of acute diarrhea cases seen in general practice was estimated at **135 cases per 100,000 inhabitants (95% CI [119 ; 151])**. This rate is **stable** compared to week 2023w05 (consolidated data: 129 [116 ; 142]) and at a level of activity close to those usually observed in this period.



Complete national and regional data are available on the last page of this bulletin.

The purpose of acute diarrhea surveillance is to monitor gastroenteritis outbreaks.

(*) Incidences of acute diarrhea during the 2019/2020 and 2020/2021 seasons were greatly reduced by containment and sanitary measures to control the Covid-19 pandemic. They are not included in historical comparisons.

Chickenpox

Low activity in general practice



In mainland France, last week (2023w06), the incidence rate of Chickenpox cases seen in general practice was estimated at **18 cases per 100,000 inhabitants (95% CI [11 ; 25])**. This rate is **increasing** compared to week 2023w05 (consolidated data: 9 [5 ; 13]) and at a low level of activity compared to those usually observed in this period.

Complete national and regional data are available on the last page of this bulletin.

(*) Incidences of Chickenpox cases during the 2019/2020 and 2020/2021 seasons were greatly reduced by the Covid-19 pandemic containment and health measures. They are not included in historical comparisons.

Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023

Additional data on acute respiratory infections

Modalities of ARI monitoring by the Sentinelles Network


Every year, viruses with respiratory tropism circulate in mainland France causing acute respiratory infections (ARI). These viruses are mainly influenza viruses, and other respiratory viruses such as respiratory syncytial virus (RSV), rhinovirus or metapneumovirus, but also SARS-CoV-2 (COVID-19) since 2020. They require close monitoring because they can be the cause of more or less severe epidemics.

In order to carry out this surveillance in general practice, Sentinel general practitioners have been reporting the number of cases of acute respiratory infection (ARI) seen in consultation (or teleconsultation) since March 17, 2020, according to the following definition: sudden onset of fever (or feeling of fever) and respiratory signs.

Virological surveillance is also carried out by Sentinel general practitioners and pediatricians, with the collection of a sample of ARI cases seen in consultation in order to identify the circulating viruses.


This monitoring is carried out in collaboration with Santé publique France, the National Reference Center (CNR) for respiratory infections (Pasteur Institute in Paris and Hospices Civils de Lyon), and the University of Corsica.

ARI incidence rates by age groups



Last week 2023w06, incidence rates are stable in all age groups except for the **15-64 years old** where there is an **increase** compared to the previous week.

Circulation of respiratory viruses in general practice and pediatrics




Last week (2023w06), 120 patients with ARI seen in GPs and paediatric consultations had been collected as a part of the Sentinel surveillance (salivary and nasopharyngeal samples). These samples were tested for various respiratory viruses, including SARS-CoV-2 (COVID-19) and influenza viruses. The results of the virological tests performed according to the weeks are presented in the graph above and detailed below:

- 57/120 (47.5) were positive for **influenza virus** (consolidated data for 2023w05: 82/160 (51.2%));
- 5/120 (4.2%) were positive for **SARS-CoV-2 (COVID-19)** (consolidated data in 2023w05: 5/161 (3.1%));
- 3/120 (2.5%) were positive for **respiratory syncytial virus (RSV)** (consolidated data in 2023w05: 7/161 (4.3%));
- 10/120 (8.3%) were positive for **rhinovirus (hRV)** (consolidated data in 2023w05: 17/161 (10.6%));
- 3/120 (2.5%) were positive for **metapneumovirus (hMPV)** (consolidated data in 2023w05: 10/161 (6.2%)).

Since week 2022w39 (September 26th 2022), 3,318 patients with ARI seen in GPs and paediatric consultations have been swabbed.

19 "Influenza A+Covid-19" co-infections, 1 "Influenza B+Covid-19" co-infection, 1 "Influenza A+Influenza B" co-infection and 1 "Influenza A+Covid-19+RSV" co-infection were observed. The A(H3N2)/SARS-CoV-2 co-infections were observed between weeks 2022s42 and 2023s01, the B (undetermined lineage)/SARS-CoV-2 co-infection was observed in 2023w05 and the A(H3N2)/B Victoria lineage co-infection was observed in 2023w04. The triple A(H3N2)/SARS-CoV-2/RSV co-infection was observed in 2022w50.

Evolution of ARI incidence by regions



The regional ARI incidence rates estimated for the last week 2023w06 are available on the last page.

In conclusion


Last week (2023w06), the incidence of ARI cases seen in general practice is increasing, and especially among the 15-64 years old (see graph opposite).

The increase in the ARI incidence rate is linked to the concomitant circulation of various respiratory viruses the past week (2023w06), in particular the influenza viruses (see page 4), but also the SARS-CoV-2 (Covid-19) (see page 3), the RSV (see page 5), the rhinovirus (hRV) and the metapneumovirus (hMPV) (see graph opposite).

Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023

INFLUENZA


ARI positivity rates to influenza by age groups



Number of samples and influenza positivity rates by age groups from ARI cases sampled by Sentinel physicians since 2022w39

Last week (2023w06), the influenza positivity rates of patients consulting for an ARI and sampled by Sentinel physicians were 58% and 39% respectively in the 0-14 and 15 and older age groups.

Estimated incidence of influenza cases seen in general practice




Incidence rate of influenza cases with fever > 39 observed in general practice since 2022w39 (blue) compared to previous seasons since 2014 (grey)

Last week (2023w06), the incidence rate of influenza cases seen in general practice was estimated at 121 cases per 100,000 population (95% CI [99; 143]), representing 80,430 [66,033; 94,827] new cases of influenza seen in general practice.

This rate is stable compared to the previous week (consolidated data for 2023w05 : 115 [99; 130], representing 76,173 [65,720; 86,626] new cases of influenza seen in general practice).

Circulation of influenza viruses



Cumulative influenza positivity rate by circulating influenza subtypes from ARI cases swabbed by Sentinel physicians since 2021w37

Last week (2023s06), among the 120 patients with ARI who were sampled, 57 were positive for influenza virus (47.5%). **The majority of circulating influenza viruses were type B (41/57 or 72%), and more precisely B Victoria lineage.**

Since the beginning of virological surveillance in week 2022s39 (26th September), the 1,007 confirmed influenza cases have been sampled by Sentinel general practitioners and pediatricians. They presented the characteristics below:

Clinical description of confirmed influenza cases:

- Their median age was 23 years (from 1 mois months to 89 years);
- 52% (516/1,001) were women;
- 92% (899/974) were not vaccinated against influenza;
- 15% (129/880) had risk factors for complications;
- 0,4% (3/853) were hospitalized at the end of the consultation.

Identification of influenza circulating viruses

The 1007 influenza viruses identified since the beginning of the virological surveillance were distributed as follows:

- 12.4% (125/1,007) influenza A(H1N1)pdm09 virus;
- 64.1% (646/1,007) influenza A(H3N2) virus;
- 2.0% (20/1,007) non-subtyped influenza A virus;
- 14.6% (147/1,007) influenza B of Victoria lineage virus;
- 7.1% (72/1,007) B lineage influenza virus not yet identified.

Influenza circulation by region

Last week (2023w06), among the 120 tested samples 57 (47,5%) were positive for at least one influenza virus. Influenza cases were identified in the majority of the French metropolitan regions, and in particular in:

- PACA (7/9 tested samples, or 78%),
- Bourgogne-Franche-Comté (2/3 tested samples, or 67%),
- Occitanie (8/12 tested samples, or 67%),
- Centre-Val de Loire (3/5 tested samples, or 60%),
- Bretagne (8/14 tested samples, or 57%),
- Hauts-de-France (5/9 tested samples, or 56%),
- Pays de la Loire (3/6 tested samples, or 50%).

The other French regions had a lower regional positivity rate than the French national rate (47.5%).

In conclusion

Last week (2023w06), the circulation of influenza viruses was stable compared to the previous week, but a **resumption of the influenza epidemic has been noted for the last three weeks**. This circulation extended to all French metropolitan regions. **This second epidemic wave is due to the circulation of an influenza lineage B Victoria virus, contrary to the rest of the season marked by the majority circulation of an A(H3N2) influenza virus.**

Characteristics (age, sex, risk factors, hospitalization) of confirmed influenza cases observed this season are similar to the ones observed during the past seasons.

You can find the epidemiological bulletin of Santé publique France with all the surveillance data (ambulatory and hospital) on influenza by clicking [here](#).

Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023

Covid-19

Modalities of Covid-19 monitoring by the Sentinelles Network


The Sentinel network contributes to the monitoring of the dynamics of the Covid-19 epidemic through the surveillance of cases of acute respiratory infection (ARI) seen in general practice (*defined as a fever or a feeling of fever accompanied by respiratory signs*).

For each patient presenting an ARI reported by Sentinel general practitioners, descriptive data are collected, including the results of diagnostic tests for Covid-19 (RT-PCR or antigenic test).

From this information, it is possible to estimate the number of Covid-19 cases with respiratory signs seen in general practice. These cases represent a majority share of all Covid-19 cases seen in general practice. However, it is important to note that Covid-19 cases without respiratory signs are not included in our estimates (such as those with only isolated ageusia or anosmia).

This indicator provides comparable estimates over time to monitor the dynamics of the epidemic.


ARI positivity rates to SARS-CoV-2 (Covid-19) by age groups



Number of ARI cases reported by Sentinelles physicians and SARS-CoV-2 (Covid-19) positivity rate since 2020w37

Last week (2023w06), the SARS-CoV-2 (Covid-19) positivity rates of patients consulting for ARI were 0%, 5%, and 12% respectively in the 0-14, 15-64, and 65 and older age groups.

Estimated incidence of Covid-19 cases with respiratory signs



Last week (2023w06), the incidence rate of Covid-19 cases with respiratory signs seen in general practice was estimated at 12 cases per 100,000 population (95% CI [8; 15]), corresponding to 7,966 [5,637; 10,295] new cases of Covid-19 with respiratory signs seen in general practice.

This rate is stable compared to those in recent weeks (consolidated data for 2023w05: 7 [5; 9], representing 4,574 [3,170; 5,978] new cases of Covid-19 with respiratory signs seen in general practice).

Description of Covid-19 cases with respiratory signs

Since week 2022w39 (26th September, date of the beginning of the virological surveillance), the 325 (9.8%) Covid-19 confirmed cases with respiratory signs sampled by the Sentinel general practitioners and paediatricians had the following characteristics:

- Their median age was 51 years (range from 3 month to 91 years) and 56% (181/323) were women;
- 18% (55/306) of cases aged 12 years and older were not vaccinated against Covid-19 (no vaccine dose received);
- 32% (94/290) had risk factors for complications;
- None were hospitalized after their consultations (0/273).

In conclusion

Last week (2023w06), the incidence of Covid-19 cases with respiratory signs seen in general practice is stable compared to the previous week, and at a low activity level in comparison to the past epidemic waves (see graph opposite).


The characteristics of SARS-CoV-2 (Covid-19) positive ARI cases observed since week 2022s21 in general practice remain similar to those observed since the beginning of the pandemic.

You can find the epidemiological bulletin of Santé publique France with all the surveillance data (ambulatory and hospital) on the Covid-19 pandemic by clicking [here](#).

Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023


RSV

ARI positivity rates to RSV by age groups



Last week (2023w06), the RSV positivity rates of patients consulting for an ARI and sampled by Sentinel physicians were 0% and 4% respectively in the 0-14 and 15 and older age groups.

Estimated incidence of RSV cases with fever >39



Last week (2023w06), the incidence rate of RSV cases with fever >39 seen in general practice was stable compared to the previous weeks, and at a similar level of activity compared to past seasons at the same time.

RSV circulation by region

Last week (2023w06), among the 109 tested samples 3 (2.8%) were positive for the respiratory syncytial virus (RSV). RSV cases were identified in 3 metropolitan regions :

- Bretagne (1/14 tested samples, or 7%),
- Auvergne-Rhône-Alpes (1/17 tested samples, or 6%),
- Ile-de-France (1/23 tested samples, or 4%).

Estimated incidence of RSV cases seen in general practice

Last week (2023w06), the incidence rate of VRS cases seen in general practice was estimated at 8 cases per 100,000 population (95% CI [3; 14]), corresponding to 5,436 [1,791; 9,081] new cases of VRS seen in general practice.

This rate is stable compared to those in recent weeks (consolidated data for 2023w05 : 10 [5; 14], representing 6,456 [3,568; 9,344] new cases of VRS seen in general practice).

Description of RSV cases

Since the beginning of virological surveillance in week 2022w39 (26th September), the 328 (9.9%) confirmed RSV cases seen by Sentinel general practitioners and pediatricians had the following characteristics:

- Their median age was 22 years (from 3 months to 96 years);
- 59% (194/325) were women;
- 20% (61/301) had risk factors for complications;
- 0,3% (1/287) was hospitalized at the end of the consultation.

These characteristics are similar to those of positive RSV cases observed in past seasons in general practice (historical data : median age: 3 years; 52% women; 23% with risk factors; 0.6% hospitalized patients).

In conclusion

Last week (2023w06), the incidence of RSV cases among patients consulting for ARI in general practice was stable compared to the previous week.

The characteristics of confirmed RSV cases were similar to the characteristics of cases from previous seasons, with the exception of the observed median age, which is higher for this 2022-2023 season.

You can find all the bronchiolitis epidemiological data (outpatient and inpatient) in the Public Health France weekly bulletin by clicking [here](#).

Observed situation in general practice for the week 6 of the year 2023, from 02/06/2023 to 02/12/2023

National incidence rates over the last 3 weeks (per 100,000 inhabitants)	2023w06 (unconsolidated) Incidence rate estimations [95% confidence interval]	2023w05 Incidence rate estimations [95% confidence interval]	2023w04 Incidence rate estimations [95% confidence interval]
Acute Respiratory Infection	284 [260 ; 308]	231 [213 ; 249]	183 [168 ; 198]
Acute diarrhea	135 [119 ; 151]	129 [116 ; 142]	125 [112 ; 138]
Chickenpox	18 [11 ; 25]	9 [5 ; 13]	9 [6 ; 12]

Regional incidence rates for the week 2023w06 (per 100,000 inhabitants)	Acute Respiratory Infection Incidence rate estimations [95% confidence interval]	Acute diarrhea Incidence rate estimations [95% confidence interval]	Chickenpox Incidence rate estimations [95% confidence interval]
Auvergne-Rhône-Alpes	291 [221 ; 361]	92 [55 ; 129]	6 [0 ; 15]
Bourgogne-Franche-Comté	233 [95 ; 371]	137 [14 ; 260]	46 [3 ; 89]
Bretagne	97 [59 ; 135]	78 [42 ; 114]	6 [0 ; 18]
Centre-Val de Loire	146 [93 ; 199]	239 [68 ; 410]	5 [0 ; 13]
Corse	220 [121 ; 319]	51 [0 ; 104]	53 [0 ; 117]
Grand Est	505 [387 ; 623]	224 [138 ; 310]	9 [0 ; 23]
Hauts-de-France	184 [118 ; 250]	170 [102 ; 238]	11 [0 ; 25]
Ile-de-France	259 [192 ; 326]	77 [50 ; 104]	10 [3 ; 17]
Normandie	96 [40 ; 152]	91 [27 ; 155]	22 [0 ; 83]
Nouvelle-Aquitaine	178 [115 ; 241]	98 [54 ; 142]	11 [0 ; 26]
Occitanie	338 [257 ; 419]	112 [61 ; 163]	2 [0 ; 6]
Pays de la Loire	82 [35 ; 129]	211 [120 ; 302]	3 [0 ; 10]
Provence-Alpes-Côte d'Azur	613 [453 ; 773]	152 [71 ; 233]	40 [5 ; 75]

French Sentinel network

Pierre Louis Institute of Epidemiology and Public Health

UMR-S 1136 (Inserm - Sorbonne Université)

Phone : +33 144 738 435 | E-mail : sentinelles@upmc.fr

Since 1984, the "réseau Sentinelles" or Sentinelles network has been a research and health monitoring network in primary care (general medicine and paediatrics) in metropolitan France. The participation of physicians is voluntary. Currently, 580 physicians participate in the continuous surveillance activity (531 general practitioners and 49 paediatricians), allowing the production of weekly epidemiological reports.

Heads of Sentinel Network : Olivier Steichen, Thierry Blanchon

Publication : Yves Dorléans

Information system & biostatistics : Clément Turbelin

Monitoring manager : Marion Debin, Caroline Guerrisi

Regional branches	Heads
Auvergne-Rhône-Alpes, Bourgogne-Franche-Comté	Marianne Sarazin
Centre-Val de Loire, Pays de la Loire	Thierry Prazuck
Corse	Alessandra Falchi
PACA	David Darmon
Grand Est	Daouda Niaré
Ile-de-France, Hauts-de-France	Mathilde François
Bretagne, Normandie	Marie Pouquet
Nouvelle-Aquitaine, Occitanie	Maryse Lapeyre-Mestre

[See all the team](#)

Need more information?

Visit our website :

<https://www.sentiweb.fr/>

You are a french GP ?

Contact us

<https://www.sentiweb.fr/france/fr/inscrire>

Envie de participer à la veille sanitaire ?

covidnet.fr

Devenez acteur de la surveillance du Covid-19 !

Pas besoin d'être malade pour participer !

Inscrivez-vous sur <https://www.covidnet.fr/>