Observed situation in primary care


Epidemiological surveillance bulletin for the week 19 of the year 2025, from 05/05/2025 to 05/11/2025


Sentinelles

Acute Respiratory Infection (ARI)

Covid-19, Influenza and other respiratory viruses

Low activity in general practice





Incidence rates and comparison with historical data

In mainland France, last week (2025w19), the incidence rate of acute respiratory infection (ARI) cases consulting in general practice was estimated at **68 cases per 100,000 population (95% CI [56; 81])**.

Subject to future data consolidation, this rate is **stable** compared to the previous week and corresponds to a **similar activity level** compared to those usually observed at this time of the year (consolidated data for 2025w18: 61 [51; 70]).


ARI are caused by a variety of respiratory viruses including SARS-CoV-2 (Covid-19), influenza viruses, and other respiratory viruses such as RSV, rhinovirus and metapneumovirus. The purpose of ARI surveillance is to monitor outbreaks of these virus.


You can find the french "Santé Publique France epidemiological bulletin" with all surveillance data (ambulatory and hospital) on ARI by clicking here.

Data sources: Sentinelles, Electronic Medical Records (EMR) IQVIA

Acute diarrhea

Low activity in general practice





Incidence rates and comparison with historical data

In mainland France, last week (2025w19), the incidence rate of acute diarrhea cases seen in general practice was estimated at **59 cases per 100,000 population** (95% CI [47; 71]).


Subject to future data consolidation, this rate is **stable** compared to the previous week and corresponds to a **similar activity level** compared to those usually observed at this time of the year (consolidated data for 2025w18: 58 [46; 69]).


The purpose of acute diarrhea surveillance is to monitor gastroenteritis outbreaks.

Data sources: Sentinelles, Electronic Medical Records (EMR) IQVIA

Chickenpox

Low activity in general practice






Incidence rates and comparison with historical data


In mainland France, last week (2025w19), the incidence rate of Chickenpox cases seen in general practice was estimated at 7 cases per 100,000 population (95% CI [2; 12]).

Subject to future data consolidation, this rate is **stable** compared to the previous week and corresponds to a **low activity level** compared to those usually observed at this time of the year (consolidated data for 2025w18: 7 [4; 11]).

Data sources: Sentinelles, Electronic Medical Records (EMR) IQVIA







Data sources: Sentinelles Network and EMR (Electronic Medical Records) IQVIA


Activity levels

- Low activity
- Moderate activity
- High activity

For the three indicators, the blue curve corresponds to the change in the incidence rate per 100,000 population for the current year. For ARI, previous years (since 2020) are shown with the grey curves. For acute diarrhea and chickenpox, the distribution of weekly incidence rates for the previous years is shown in grayed colour, with quartiles in darker and minimum/maximum values in lighter and minimum/maximum values in lighter and minimum/maximum values in lighter because the current trends to be compared with historical data. The value of the last point and its confidence interval are shown at the top of each graph. Different scales are used for different indicators.

Sentinelles

Incidence rates of Covid-19 casesDecreasing activity stable and at a low level




National ARI incidence rate due to Covid-19 and comparison with historical data

Last week (2025w19), the incidence rate of **Covid-19** cases seen in general practice among patients consulting for an ARI was estimated at **3 cases per 100,000 population** (95% CI [2; 5]), corresponding to 2 313 [1 042; 3 584] new cases.

Subject to future data consolidation, this rate is **decreasing** compared to the previous week (consolidated data for 2025w18: 9 [6; 12]).

Incidence rates of Covid-19 cases by age groups



Last week (2025w19), the incidence rates of **Covid-19** cases seen in general practice among patients consulting for an ARI were estimated at:

- **0-14 years**: 0 cases per 100,000 population;
- **15-64 years**: 3 cases per 100,000 population (95% CI [1; 5]), corresponding to 1,318 [525; 2,112] new cases;
- **65 years and above**: 7 cases per 100,000 population (95% CI [1; 13]), corresponding to 995 [146; 1,844] new cases.

Subject to future data consolidation, these rates are decreasing in the 65 years and above age group, and stable in the other age groups, compared to those of the previous week.

Description of Covid-19 cases presenting ARI seen in general practice

Since week 2025w16, the 67 Covid-19 described cases with an ARI had the following characteristics:

- Median age: 63 years (range from 15 years to 93 years);
- Male/female sex-ratio: 0.72 (28/39);
- **Risk factors**: 34% (23/67) of the patients had risk factors for complications;
- **Hospitalization**: 3% (2/67) of the patients were hospitalized after the consultation.

Data source: Sentinelles

In conclusion

Last week (2025w19), subject to future data consolidation, the incidence of **Covid-19** cases seen in general practice among patients consulting for an ARI was **decreasing** compared to the previous week and remained at a **low level of activity**.

Further information is available on the "Santé publique France" website

Data source: Sentinelles

General organization and partners

Sentinelles

Surveillance organisation

Under the aegis of Santé publique France, surveillance in general practice in mainland France is moving towards the integration and joint analysis of data from different networks.

The epidemiological surveillance data published in this bulletin come from several complementary networks of general physicians:

- The Sentinelles network, coordinated by the Institut Pierre Louis of Epidemiology and Public Health (iPLESP) under the supervision of Sorbonne University and Inserm;
- and the EMR (Electronic Medical Records) database. managed by IQVIA.

During the enhanced respiratory infection surveillance season (September to April), data are also collected from physicians in the network coordinated by the general medicine departments of the University of Rouen and the Côte d'Azur University.

All these collected data are analysed jointly. They provide more reliable on a finer geographical scale, while limiting consolidation from one week to the next.

Current monitoring concerns nine health indicators with three of them being published each week in this bulletin:

You can find more information about the organization of this surveillance, the number of participating physicians, the methods used, scientific publications and partnerships on the Sentinelles network website: www.sentiweb.fr

Information and contacts

The Sentinelles team is composed of epidemiologists. statisticans, physicians, IT specialists and technicians.

Head of the Sentinelles network Olivier Steichen, Thierry Blanchon

IT Biostatistics Clément Turbelin

Epidemiological Surveillance and Studies Marion Debin

Publication Yves Dorléans

CONTACT US

- 06 64 84 66 62
- sentinelles@upmc.fr
- **IPLESP UMR-S 1136** Faculté de Santé Sorbonne Université Site Saint-Antoine, BC 2908 27, rue Chaligny 75571 Paris Cedex 12

Partners and supervisory bodies

Partners

























Supervisory bodies of Sentinelles network







French General Practionner or Paediatrician?



Get involved in research and health monitoring in primary care by joining the Sentinelles network (become a Sentinelles doctor)!

THERE IS ALSO GENERAL POPULATION MONITORING



Join the participatory cohort for monitoring Covid-19 and influenza by registering at https://www.grippenet.fr

You don't need to be a healthcare professional to take part!